

A new flea beetle genus from sub-Saharan Africa, and taxonomic remarks on the *Blepharida* genera group in the Afrotropical region (Chrysomelidae, Galerucinae, Alticini)

Paola D'Alessandro¹, Maurizio Biondi¹

¹ Department of Health, Life and Environmental Sciences, University of L'Aquila, 67100 L'Aquila, Italy

Corresponding author: Paola D'Alessandro (paola.dalessandro@univaq.it)

Abstract

Cladocera fulvipennis Jacoby is transferred to *Afrotropicaltica* **gen. nov.**, here described. The new genus belongs to the group of Afrotropical genera traditionally attributed to the *Blepharida* group: *Blepharidina* Bechyné, *Calotheca* Heyden, *Diamphidia* Gerstaecker, *Polyclada* Chevrolat, and *Xanthophysca* Fairmaire. A diagnostic key to these genera, and the list of species currently attributed to *Diamphidia* and *Polyclada* are provided based on type material and original species descriptions. Species and genera whose taxonomic position needs a revision were identified. Finally, the following synonymy is proposed: *Cladocera fulvipennis* Jacoby, 1895 = *Blepharida favareli* Achard, 1922, **syn. nov.**

Key words: *Afrotropicaltica*, *Blepharidina*, *Calotheca*, *Cladocera fulvipennis*, *Diamphidia*, new synonymy, *Polyclada*, *Xanthophysca*



Academic editor:

Alexander Konstantinov

Received: 18 October 2024

Accepted: 5 December 2024

Published: 20 February 2025

ZooBank: <https://zoobank.org/B4C6A019-EA05-4955-BCB0-159CE81B345C>

Citation: D'Alessandro P, Biondi M (2025) A new flea beetle genus from sub-Saharan Africa, and taxonomic remarks on the *Blepharida* genera group in the Afrotropical region (Chrysomelidae, Galerucinae, Alticini). ZooKeys 1228: 139–159. <https://doi.org/10.3897/zookeys.1228.139654>

Copyright: ©

Paola D'Alessandro & Maurizio Biondi.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

Introduction

The term *Blepharida* group refers to a group of approximately 21 genera from the Afrotropical, Nearctic, Neotropical, and Oriental regions (Furth 1992, 1998, Medvedev 1999; Furth and Lee 2000; Biondi et al. 2017; Calcetas and Staines 2024). Various authors discussed similarities and affinities between the genera based on adult and larval morphology, molecular markers, and ecology, and established new nomenclatural acts, generally focusing on one subset of taxa: Furth (1992, 1998), Medvedev (1999), Furth and Lee (2000), Becerra (2004), Takizawa (2005), Chaboo et al. (2007), Lee and Cheng (2007), Prathapan and Chaboo (2011), Biondi et al. (2017). Furth and Lee (2000) provided a morphological synthesis of the group based on adult characters (tarsal claws, procoxal cavities, head, pronotum, hind femora, eye, proepimeron, and metatibia) and larval characters (antenna, mandible, labrum, stemmata, endocarina, coronal suture, frontal suture), but highlighted that some characters are not shared by all the genera.

The Afrotropical region hosts five genera traditionally considered to belong to the *Blepharida* group, sensu Furth and Lee (2000): *Blepharidina* Bechyné, 1968; *Calotheca* Heyden, 1887; *Diamphidia* Gerstaecker, 1855; *Polyclada* Chevrolat, 1837, and *Xanthophysca* Fairmaire, 1901 (Biondi and D'Alessandro 2010, 2012; Biondi et al. 2017, 2022). Genus *Blepharidina* has recently been the object of a deep

taxonomic revision (Biondi et al. 2017, 2019; D'Alessandro et al. 2018a, 2019). It includes 32 species, of which 12 are in the subgenus *Afroblepharida* Biondi & D'Alessandro, 2017, and 20 in the subgenus *Blepharidina* s. str. The genus mainly occurs in the intertropical area of Africa, with *Blepharidina* primarily distributed in the south and *Afroblepharida* largely in the central-eastern area, including Socotra Island, with extensions towards the north and in the west. Moreover, *Blepharidina* s. str. species generally occur in mesic environments, while *Blepharidina* (*Afroblepharida*) species are generally associated with more xeric conditions (D'Alessandro et al. 2018b; Iannella et al. 2021). *Calotheca* comprises 37 described species (Biondi et al. 2017; D'Alessandro et al. 2020, 2021, 2022, 2023a, 2023b). Some species groups have been revised recently, while others are still under investigation by the authors. The genus is widespread in sub-Saharan Africa, and particularly common in the eastern and southern parts of its distribution range, with limited extensions into Israel and the Arabian Peninsula (Iannella et al. 2021). *Calotheca* species are generally associated with savannas and forests, or with the typical South African vegetation, such as Albany thicket and Fynbos (D'Alessandro et al. 2018b; Iannella et al. 2021; Biondi et al. 2024). Many records reported *Searsia* species (Anacardiaceae) as the primary host plants (Koch 1958; Furth and Young 1988; MB, pers. data; E. Grobbelaar, pers. comm. 10 September 2024). *Polyclada* occurs in sub-Saharan Africa and the Arabian Peninsula with ~ 15 species (Biondi and D'Alessandro 2010, 2012; D'Alessandro and Biondi 2023). Species are associated with Anacardiaceae and Burseraceae and are found in various woodland and savannah ecosystems (Chaboo et al. 2007; Prathapan and Chaboo 2011; Biondi et al. 2022). *Diamphidia* comprises ~ 17 species, widespread in central, eastern, and southern-western Africa (Biondi and D'Alessandro 2010, 2012) and associated with *Commiphora* shrubs and trees (Burseraceae) (Chaboo et al. 2007; Biondi and D'Alessandro 2012). *Xanthophysca* is endemic to Madagascar, and currently includes 5–7 species (Biondi and D'Alessandro 2010, 2012; Biondi et al. 2017).

Diamphidia, *Polyclada*, and *Xanthophysca* have not been revised recently. *Polyclada* and *Diamphidia* species, previously described also under the generic names *Cladocera* Hope, 1840 and/or *Cladotelia* Kolbe, 1894, created a certain confusion in the generic name usage and, regarding *Polyclada*, also the genus author (Baly 1861; Chapuis 1875; Achard 1922; Laboissière 1941, 1942; Bryant 1942). In addition, discordances in the identification and description of key diagnostic characters for the genera *Diamphidia* and *Polyclada* have led to uncertainty in the attribution of some species and/or the transferring from one genus to the other (Baly 1861; Chapuis 1875; Achard 1922; Laboissière 1941, 1942; Bryant 1942).

In this paper, we establish the synonymy *Cladocera fulvipennis* Jacoby, 1895 = *Blepharida favareli* Achard, 1922, syn. nov., designate the lectotype of *Blepharida favareli*, and provide a reassessment of the taxonomic position of *Cladocera fulvipennis* by transferring it to *Afrotropicaltica* gen. nov. This new flea beetle genus has been established after the revision of the diagnostic characters of the Afrotropical genera traditionally attributed to the *Blepharida* group, specifically the adult morphological characters. Based on this revision, a diagnostic key to the genera is also provided. Since *Blepharidina* and most *Calotheca* species were revised and listed in recent papers (Biondi et al. 2017; D'Alessandro et al. 2018a, 2019, 2020, 2021, 2022, 2023a, b), particular attention is paid to *Diamphidia*, *Polyclada*, and *Xanthophysca*. The list of species attributed to these three genera is provided. Since the most recent taxonomic literature

does not agree on some synonymies and/or the status of some taxa, possible synonyms are listed as separated species. Species whose generic attribution needs further investigation are reported. *Cladocera* and *Cladotelia* are currently considered as synonyms of *Polyclada* (Biondi and D'Alessandro 2012).

Materials and methods

Material examined consisted of dried pinned specimens preserved in the depositories listed in the “Abbreviations” section; abbreviations followed the list on the website The Insect and Spider Collections of the World (Evenhuis 2021). Species are attributed to the genera *Diamphidia*, *Polyclada*, and *Xanthophysca* focusing on type material and original species descriptions. Species whose taxonomic position at the genus level needs a revision are listed separately. Exact label data are cited for all type specimens; a double slash (//) divided the data on different labels and a single slash (/) divided the data in different rows. Information included in square brackets has been added to the label data using the Google Earth website for coordinates and geographic information. Geographic coordinates for the localities were reported in Degrees and Decimal Minutes (DDM) format using the WGS84 datum. [?] refers to undetectable or doubtful locality or illegible handwriting. Specimens were examined, measured, and dissected using a Leica M205C stereomicroscope. Photographs were taken using a Leica DMC5400 camera and compiled using Zerene Stacker software, v. 1.04. Scanning electron micrographs were taken using a Hitachi TM-1000. Terminology for genitalia follows Döberl (1986), Schmitt and Uhl (2015: fig. 1), and D'Alessandro et al. (2016: figs 10E, 11E). Terminology for the metafemoral spring or metafemoral extensor tendon (Nadein and Betz 2016) follows Furth (1982).

Abbreviations

Collections and depositories

BAQ	Italy, University of L'Aquila, Collection of M. Biondi;
MCZC	USA, Massachusetts, Cambridge, Harvard University, Museum of Comparative Zoology;
MNHN	France, Paris, Muséum National d'Histoire Naturelle;
MSNG	Italy, Genova, Museo Civico di Storia Naturale di Genova;
NHMUK	United Kingdom, London, The Natural History Museum;
NMPC	Czech Republic, Prague, National Museum (Natural History);
RMCA	Belgium, Tervuren, Musée Royal de l'Afrique Centrale.

Biometrics

LA	numerical sequence from base to apex of each antennomere, proportional to the length of the first antennomere;
LAED	length of median lobe of the aedeagus;
LAN	length of antennae;
LB	total body length (from apical margin of head to apex of elytra);
LE	length of elytra;
LF	maximum length of hind femora;

LP	medial length of pronotum;
LSP	maximum length of spermatheca, including ductus;
WE	maximum width of elytra combined;
WF	maximum width of hind femora;
WP	maximum width of pronotum.

Taxonomic account

Afrotropicaltica gen. nov.

<https://zoobank.org/011F6C24-9303-4535-9952-09533542B544>

Fig. 1

Description. Body subelliptical-elongate in dorsal view, with subparallel sides (Fig. 1A, B), distinctly convex in lateral view. Dorsal surface glabrous, bicoloured in the only known species, with yellowish head and pronotum, and reddish brown elytra. Head (Fig. 1E) with frontal grooves, frontal carina, and frontal calli barely distinguishable; surface smooth or micropunctate, with rounded punctures approx. as large as the supraorbital setiferous pore; puncture absent on the middle of front and vertex; eyes large, ovoidal; minimum distance between eyes on vertex, as wide as $2 \times$ the interantennal space; interantennal space prominent compared to the clypeus, as wide as $\sim 2/3$ the length of the first antennomere; maxillary palpi three-articulated, and labial palpi bi-articulated, both with slender, subcylindrical segments. Antennae (Fig. 1A, B) with 11 antennomeres, filiform, longer than $1/2$ the body length in both sexes, but slightly shorter in female; antennomere II slightly shorter than $1/2$ of antennomere I and as long as $1/2$ of antennomere III; antennomere IV as long as (female) or slightly longer (male) than antennomere III; antennomeres V–IX generally slightly longer than antennomere I (male) or approximately as long as antennomere I (female); antennomere X slightly shorter than antennomere I, especially in female; antennomere XI distinctly longer than antennomere I. Prothorax distinctly depressed dorsally (not subcylindrical). Pronotum (Fig. 1D) subrectangular, slightly convergent anteriorly, with curved, moderately expanded lateral margins, clearly visible in dorsal view; basal margin moderately arched; surface smooth to micropunctate without any groove or depression; anterior angles barely prominent; posterior angles widely obtuse; punctation rounded, quite sparse and irregularly distributed. Scutellum subtriangular. Elytra (Fig. 1A, B, D) subparallel, with base as wide as pronotal base; lateral margins finely bordered, barely visible in dorsal view; punctation rounded, variably arranged in numerous rows or bands, or almost confused. Epipleurae oblique, clearly visible in lateral view. Procoxal cavities open posteriorly; prosternum approx. as wide as the procoxal cavities; intercoxal prosternal process quite narrow; mesosternum slightly wider than prosternum, and distinctly wider than mesocoxal cavities; metasternum wider than mesosternum, twice the width of the metacoxal cavities; first abdominal ventrite as wide as metacoxal cavities. Posterior femora elliptical-elongate, moderately swollen; dorsal margin of middle and hind tibiae with distinct ciliate emargination, acute apically; hind tibiae straight in dorsal view; apical spur of hind tibiae simple; third visible metatarsomere deeply incised; fourth visible tarsomere of metatarsus simple, not swollen in both male and female; claws appendiculate. Metafemoral extensor tendon (Fig. 1H) with extended arm (ea: Fig. 1H) as long as $\sim 1/2$ the dorsal lobe (dl: Fig. 1H); dorsal-basal angle (dba: Fig. 1H)




Figure 1. *Afrotropicaltica fulvipennis* (Jacoby) comb. nov. **A** lectotype of *Blepharida favareli* Achard, male, Ogooué, Sam-Kita, habitus **B** holotype of *Cladocera fulvipennis* Jacoby, male, habitus **C** distribution **D** head, pronotum, and base of elytra, male, Cameroun, Victoria **E** head, Republique du Congo, P.N. d'Odzala, Mbandza **F** median lobe of the aedeagus, from left to right in ventral, dorsal, and lateral view, Republique du Congo, P.N. d'Odzala, Mbandza **G** spermatheca, paralectotype of *Blepharida favareli* Achard, Democratic Republic of the Congo, Eala **H** metafemoral extensor tendon, Cameroun, Victoria. Abbreviations: am: apical margin; bav: basal angle of the ventral lobe; dba: dorsal-basal angle; dl: dorsal lobe; dmrv: dorsal margin of the ventral lobe; ea: extended arm; vba: ventral-basal angle; vl: ventral lobe. Scale bars: 3 mm (**A**, **B**); 1 mm (**D**, **E**, **F**, **G**); 500 μ m (**H**).

approx. right angled, moderately prominent apically; ventral-basal angle (vba: Fig. 1H) distinctly obtuse and moderately rounded; apical margin of the tendon (am: Fig. 1H) slightly C-shaped; dorsal margin of ventral lobe (dmv: Fig. 1H) oblique; basal angle of ventral lobe (bav: Fig. 1H) quite close to the dorsal-basal angle of the tendon. Median lobe of the aedeagus (Fig. 1F) mostly subparallel in ventral view, with truncated apex, and curved in lateral view; dorsal ligula formed by a wider central lobe, and two lateral lobes. Spermatheca (Fig. 1G) with subovate basal part, narrowing towards ductus attach; distal part thin, without a distinct collum.

Type species. *Cladocera fulvipennis* Jacoby, 1895.

Etymology. The name of the new genus refers to flea beetle with strictly Afrotropical distribution. Gender feminine.

Distribution. Cameroon, Central African Republic, Democratic Republic of the Congo, Gabon, Ivory Coast, Kenya, Republic of South Africa, Togo, Uganda (Fig. 1C).

***Afrotropicaltica fulvipennis* (Jacoby, 1895), comb. nov.**

Fig. 1

Cladocera fulvipennis Jacoby, 1895: 179.

= *Blepharida favareli* Achard, 1922: 8, syn. nov.

Blepharida fulvipennis (Jacoby, 1895): Achard 1922: 9, Bryant 1942: 161.

Diamphidia fulvipennis (Jacoby, 1895): Bechyné 1960: 111, Scherer 1969: 371.

Diamphidia fulvipennis favareli (Achard, 1922): Bechyné 1960: 111, Scherer 1962: 78, Scherer 1972: 16.

Diamphidia favareli (Achard, 1922): Laboissière 1941: 240, Biondi et al. 2017: 112.

Comments. Achard (1922) described *Blepharida favareli* and transferred *Cladocera fulvipennis* Jacoby to the genus *Blepharida*, due to the similarities with *B. favareli*. Bryant (1942) transferred *Cladocera fulvipennis* to *Blepharida*, as already proposed by Achard (1922). Bechyné (1960) established the combination *Diamphidia fulvipennis* (Jacoby, 1895) and considered the species described by Achard, *Blepharida favareli*, as a geographic form of *D. fulvipennis*. Scherer (1969) erroneously reported *Diamphidia fulvipennis* (Jacoby, 1895) as *Diamphidia fulvipennis* Jacoby, 1893. We formalize here the synonymy between *Cladocera fulvipennis* Jacoby, 1895 and *Blepharida favareli* Achard, 1922.

Type material examined. **Holotype** ♂ of *Cladocera fulvipennis*: "Type H.T // Togo / Africa // 42 // Type // Jacoby Coll. / 1909–28a. // *Cladocera / fulvipennis* / Jac. / Type" [8° 31.995'N 1° 5.853'E] (NHMUK). MCZC hosts the following specimen: "Type / 18327 // Boma / M. Tschoffen // Jacoby 2nd / Coll. // fulvipennis Jac." [Democratic Republic of the Congo, Boma, 5° 50.762'S 13° 4.296'E 5°52'32"S, 13°02'00"E] (photos available at <https://mczbase.mcz.harvard.edu/guid/MCZ:Ent:18527>). Lectotype ♂ of *Blepharida favareli* Achard, here designated: "Type // B. Favareli / Type / J. Achard det. // Ogooué / Sam-Kita" [Gabon, Samkita, 0° 26.516'S 10° 27.247'E] (NMPC). Paralectotype ♀ of *Blepharida favareli* Achard, here designated: "Type // B. Favareli / Type / J. Achard det. // Fort-Sibut / Oubanghi-Chari" [Central African Republic, Fort-Sibut, 5°43.57140'N 019°05.11590'E] (NMPC). Paralectotype ♀ of *Blepharida favareli* Achard, here designated: "Type // B. Favareli / Type / J. Achard det. // Musée du Congo / Eala / IX 1912 / R. Mayré" [Democratic Republic of Congo, Eala, 0° 2.441'N 18° 20.149'E] (NMPC).

Other material. CAMEROON • 16 specs, Victoria [= Limbé 4°1.917'N, 9°11.122'E], Jun.–Jul.1902, L. Fea leg. (MSNG). CENTRAL AFRICAN REPUBLIC • 1♀, Bamingui-Bangoran Pr., 45 km SSW Bamingui [7°16.400'N, 19°55.580'E], 450 m, 13–15 May 2009, A. Kudrna Jr leg. (BAQ). DEMOCRATIC REPUBLIC OF CONGO • 1 spec., Boma [5°50.762'S, 13°4.296'E], M. Tschoffen leg., Jacoby Coll. 1909-28a (NHMUK); 2 specs, ibid (RMCA); 1 spec., Eala [0°2.441'N, 18°20.149'E], Oct. 1935, J. Ghesquière leg. (RMCA); 1 spec., P.N. d'Odzala [1°16.167'N, 14°52.299'E], Mbandza, Nov. 1992, G. Carpaneto leg. (BAQ); 1 spec., Ekoiongouma, N1.20279 E17.88032 [1°12.16740' N, 17°52.81920' E], 341 m, 14 Feb. 2022, canopy light trap (-10 m), C.N. Hackfort & A. Tsoumou leg. (NHMUK); 1 spec., Lokandu [2°31.228'S, 25°44.566'E], Mar.1939, [Capt.] Marée leg. (NHMUK); 13 specs, ibid (RMCA); 1 spec., ibid, 1939 (RMCA); 3 specs, Equateur, Flandria [0°22.775'S, 19°5.500'E], 27 Oct.1932, R.P. Hulstaert leg. (RMCA); 1 spec., ibid, 31 Mar. 1932; 1 spec., Flandria, 15 Mar. 1932; 1 spec., ibid, 1931; 1 spec., Cataractes de Luozi [Luozi 4°54.711'S, 14°9.842'E], Dec. 1898, E. Luja leg. (RMCA); 1 spec., Bas-Congo, Lemfu [5°21.206'S, 15°15.709'E], Jun. 1945, Rév. P.L. De Reir leg. (RMCA); 1 spec., Tshuapa, Bokuma [0°40.648'S, 21°1.305'E], Dec. 1951, R.P. Lootens leg. (RMCA); 10 specs, Kisantu [5°7.473'S, 15°7.749'E], P. Goossens leg. (RMCA); 1 spec., ibid, ex col. Seeldrayers; 3 specs, ibid, 1905, col. Clavareau; 1 spec., Uelé, Dingila [3°38.633'N, 26°3.515'E], Jun. 1933, J.V. Leroy leg. (RMCA); 1 spec., Uelé, Paulis [= Isiro 2°46.212'N, 27°37.242'E], 1947, Abbeloos leg. (RMCA); 1 spec., Uelé, Bambessa [3°27.170'N, 25°41.579'E], 15 Sept. 1933, Lefèvre leg. (RMCA); 1 spec., Stanleyville [= Kisangani 0°30.542'N, 25°13.409'E], 1924, J. Ghesquière leg. (RMCA); 3 specs, Dima [3°16.47444'S, 17°28.99290'E], 14 Sept. 1908, A. Koller leg. (RMCA); 1 spec., Congo da Lemba [Lemba 4°24.598'S, 15°20.277'E], Jan. 1913, R. Mayné leg. (RMCA); 1 spec., ibid, 1–15 Apr. 1913; 7 specs, ibid, oct.–Dec.1911; 1 spec., Mongbwalu (Kilo) [1°55.640'N, 30°2.904'E], 1939, Mme Scheitz leg. (RMCA); 1 spec., Kilo [Kilo-Etat 1°49.738'N, 30°9.391'E], Dr. Abetti leg. (RMCA); 1 spec., Ganda Sundi [4°52.000'S, 12°52.000'E], [Comte J.] de Briey leg. (RMCA); 1 spec., Mahagi-Niarembe [between Mahagi and Niarembe 2°16.917'N, 31°3.371'E], 1935, Ch. Scops leg. (RMCA); 3 specs, Sud Ogowé, 'Ngomo [4°57.378'S, 23°37.289'E], [H.A.] Junod leg. (RMCA); 2 specs, Bas Congo, Maydi [5°11.958'S, 15°9.195'E], 1945, Rév. P. Van Eylen leg. (RMCA); 21 specs, ibid; 28 specs, ibid, 1942; 2 specs, ibid, 1943; 3 specs, Kivu, Kavumu à Kabunga, km 82 (Mingazi) [2°1.870'S, 28°30.949'E], Apr.–Jul.1951, H. Bomans leg. (RMCA); 1 spec., ibid, Nov.–Dec. 1951; 1 spec., ibid, 1951; 1 spec., Bas-Uele, Buta [2°47.787'N, 24°44.473'E], 1926, Fr. Joseph leg. (RMCA); 1 spec., Mayumbe, Makaia N'Tete [Mayumbe 2°30.000'N, 27°37.000'E], 24 Nov. 1915, R. Mayné leg. (RMCA); 1 spec., Busira [0°5.233'S, 19°55.086'E], 18 Oct.1905, Waelbroeck leg. (RMCA); 1 spec., Equateur, Bohuma [1°8.099'N, 30°13.892'E], 1938, R.P. Hulstaert leg. (RMCA); 1 spec., Ilenge [1°58.940'S, 19°52.474'E], 30 Jan. 1913, R. Mayné leg. (RMCA). GABON • 1♀, Gabon Français [0°59.062'S, 11°54.238'E] (NMPC). IVORY COAST • 1 spec., Andé, Bongouanou [Bongouanou 6°39.047'N, 4°11.247'W], Mar. 1962, J. Decelle leg. (RMCA); 1 spec., 1171 m, Mt Tonkouï Peak, 07°27'15.2"N, 07°38'12.5"W [7°27.230'N, 7°38.148'W], 1–8 Nov. 2015, general collecting, M. Aristophanous, P. Moretto, E. Ruzzier leg. (NHMUK). KENYA • 1 spec., Brit. E. Africa, Nyangori, N Kavirondo [0°41.195'S, 35°0.679'E], 4800 ft, 18–19 May 1911, S.A. Neave leg. (NHMUK). UGANDA • 1 spec., Mawakota [Mawokota region 0°36.800'S, 30°12.695'E], Nov. 1931, [V.G.L.] van Someren leg. (NHMUK); 1 spec., Boy [Bay (?)], Entebbe [0°3.290'N,

32°28.216'E], 2800-4000 ft, forest within four miles of Kitabi Hill, May–Jun. 1913, C.A. Wiggins leg. (NHMUK); 1 spec., Entebbe, 14 Aug. 1911, C.C. Gowdey leg. (NHMUK); 1 spec., Prot. Entebbe, 12–20 Jan. 1912, S.A. Neave leg. (RMCA); 1 spec., Kampala [0°19.253'N, 32°35.055'E], 1–10 Jan. 1918, C.C. Gowdey leg. (NHMUK); 1 spec., Shores of L. Isolt, or Wamala [Wamala 0°7.799'N, 31°38.115'E], 3800 ft, 7–8 Jan. 1912, S.A. Neave leg. (NHMUK).

Redescription. Body subelliptical-elongate in dorsal view with parallel sides (Fig. 1A, B), distinctly convex in lateral view; total length of the body (LB) = 8.10 ± 0.91 mm ($6.50 \leq LB \leq 9.70$ mm) in male, and 8.77 ± 0.65 mm ($7.52 \leq LB \leq 9.51$ mm) in female; maximum pronotal width in the middle: WP = 3.43 ± 0.29 mm ($2.86 \leq WP \leq 3.78$ mm) in male, and 3.71 ± 0.25 mm ($3.30 \leq WP \leq 3.98$ mm) in female; maximum width of elytra in the middle: WE = 4.37 ± 0.37 mm ($3.69 \leq WE \leq 4.85$ mm) in male, and 4.94 ± 0.42 mm ($4.46 \leq WE \leq 5.72$ mm) in female; WE/WP = 1.27 ± 0.04 ($1.22 \leq WE/WP \leq 1.33$) in male, and 1.33 ± 0.05 ($1.29 \leq WE/WP \leq 1.44$) in female.

Head and pronotum yellowish (Fig. 1A, B, D); elytra reddish brown; scutellum paler than elytra, often of the same colour as the pronotum or slightly darker; underside yellowish to pale brown; mouthparts yellowish, with black distal part of mandibles; antennae and legs with two colour patterns: both antennae and legs rather homogenous, yellowish to reddish brown, at most with slightly paler tarsi; antennal segments 1–4 yellowish and 5–11 blackish, and legs mostly blackish with the exception of femora, paler in their proximal part. Head (Fig. 1E) with smooth to micropunctate surface and sparse, rounded punctures, denser on vertex and near the inner ocular margins; frontal grooves and frontal calli barely distinguishable; frontal carina wide, not raised; eyes large, ovoidal; interantennal space as wide as $\sim 2/3$ the length of the first antennomere. Antennae (Fig. 1A, B) filiform, longer than $1/2$ the body length: LAN = 6.07 ± 0.44 mm ($5.43 \leq LAN \leq 6.69$ mm) in male, and 5.62 ± 0.39 mm ($5.04 \leq LAN \leq 6.31$ mm) in female, and LAN/LB = 0.75 ± 0.05 ($0.65 \leq LAN/LB \leq 0.84$) in male, and 0.64 ± 0.03 ($0.60 \leq LAN/LB \leq 0.68$) in female. LA = 100:42:83:92:117:117:117:108:100:92:125 in male, and 100:42:83:83:108:100:100:92:83:83:117 in female. Pronotum (Fig. 1D) distinctly transverse: LP = 1.79 ± 0.17 mm ($1.50 \leq LP \leq 1.94$ mm) in male, and 1.85 ± 0.12 mm ($1.65 \leq LP \leq 2.04$ mm) in female, and WP/LP = 1.91 ± 0.06 ($1.80 \leq WP/LP \leq 2.00$) in male, and 2.01 ± 0.03 ($1.95 \leq WP/LP \leq 2.05$) in female; lateral margins slightly convergent anteriorly, moderately to distinctly curved, moderately expanded, visible in dorsal view; basal margin moderately arched; surface smooth to micropunctate; main punctation formed by rounded punctures, clearly impressed, quite sparse and irregularly distributed. Scutellum subtriangular, laterally rounded. Elytra (Fig. 1A, B, D) subparallel, clearly longer than wide, jointly rounded apically. LE = 6.57 ± 0.54 mm ($5.43 \leq LE \leq 7.23$ mm) in male, and 7.21 ± 0.48 mm ($6.40 \leq LE \leq 7.86$ mm) in female; WE/LE = 0.67 ± 0.02 ($0.63 \leq WE/LE \leq 0.70$) in male, and 0.68 ± 0.02 ($0.66 \leq WE/LE \leq 0.73$) in female; LE/LP = 3.66 ± 0.12 ($3.50 \leq LE/LP \leq 3.91$) in male, and 3.90 ± 0.09 ($3.80 \leq LE/LP \leq 4.05$) in female. Lateral margins finely bordered, indistinctly visible in dorsal view; surface very finely micropunctate or microwrinkled; main punctation clearly impressed, dense, variably arranged in numerous rows or bands, or almost confused. Humeral calli distinctly raised. Macropterous. Posterior femora moderately swollen (WF/LF = 0.44 ± 0.01); apical spur of hind tibiae very short; first tarsomere of fore- and middle legs clearly enlarged in male. Median lobe of the aedeagus (Fig. 1F) in ventral view: outline mostly subparallel

with slightly sinuate margins, and narrower subapical part; apex truncated, distinctly expanded laterally; surface smooth, concave laterally and prominent medially in the apical third; in lateral view median lobe distinctly curved, with slightly sinuate outline; dorsal ligula formed by a wide central lobe, and two shorter and thinner lateral lobes, with base at apical $\sim 1/3$; LAED = 2.26 ± 0.10 mm ($2.13 \leq \text{LAED} \leq 2.43$ mm); LE/LAED = 2.90 ± 0.19 ($2.55 \leq \text{LE/LAED} \leq 3.17$). Basal part of the spermatheca (Fig. 1G) subovate, abruptly narrowing towards the ductus insertion; ductus short, thick, uncoiled, ventrally inserted; distal part of the spermatheca clearly bent towards the basal part, thin, acute apically; LSP = 0.93 ± 0.05 mm ($0.87 \leq \text{LSP} \leq 1.02$ mm); LE/LSP = 7.78 ± 0.25 ($7.33 \leq \text{LE/LSP} \leq 8.11$).

Distribution. Cameroon, Central African Republic, Democratic Republic of the Congo, Gabon, Ivory Coast, Kenya, Republic of South Africa, Togo, Uganda (Fig. 1C).

Ecological notes. Ecology, including host plant, unknown.

Key to the Afrotropical genera of the *Blepharida* group

The combination of characters identifying *Blepharidina* and *Calotheca* was based on recent revisions (Biondi et al. 2017, 2019; D'Alessandro et al. 2018a, 2019, 2020, 2021, 2022, 2023a, 2023b). Regarding *Diamphidia*, *Polyclada*, and *Xanthophysca*, characters reported in the key were derived from the species whose genus attribution was unequivocal, due to their similarities with the type species.

- 1 Antennae filiform, at most with distal segments slightly enlarged (Figs 1A, B, 2A, 3A, 6A); segment 4 approx. as long as 3 **2**
- Antennae pectinate, serrate, or with middle segments clearly subtriangular (Figs 4A, 5A); segment 4 distinctly longer than 3 **5**
- 2 Pronotal punctation not homogenous, with at least some larger and more deeply impressed punctures arranged in lines (at least one line) or dense groups (Figs 2B, C, 3B, C) **3**
- Pronotal punctation uniform in size, homogeneously or slightly irregularly distributed (Figs 1D, 4B, 5B, 6B) **4**
- 3 Pronotum with main punctation arranged in two striae that are straight, L-, or C- shaped, running from the anterior margin towards the disc (ps: Fig. 3B, C); basal longitudinal furrows or small dimples are present in some species (bf: Fig. 3B). Frontal grooves elongate, sinuate, and deeply impressed from the superior ocular margin to the interantennal space (fg: Fig. 3E); frons surface not depressed between antennal sockets and clypeus. Extended arm of the metafemoral extensor tendon $\sim 1/3$ the length of the dorsal lobe; apical margin of the tendon clearly C-shaped (Fig. 3D) ***Calotheca* Heyden**
- Pronotum with main punctation arranged in patches, and oblique, transverse, or longitudinal lines (Fig. 2B, C). Frontal grooves, not distinguishable (Fig. 2E), or at most short and moderately impressed; frons surface depressed between antennal sockets and clypeus. Extended arm of the metafemoral extensor tendon slightly shorter than $1/2$ the length of the dorsal lobe; apical margin of the tendon moderately C-shaped (Fig. 2D) ***Blepharidina* Bechyné**
- 4 Procoxal cavities closed posteriorly. Pronotal punctation mostly uniform and dense (Fig. 6B). Frontal grooves distinctly impressed, from the antennal sockets to the upper ocular margin (Fig. 6B). Hind femur with a distinct emargination on the posterior side in both sexes, but more evi-

- dent and tooth-shaped in male (Fig. 6A). Ductus spermathecae sub-truncate, practically absent (Fig. 6E). General shape of the median lobe of the aedeagus roughly subcylindrical (Fig. 6D). Metafemoral extensor tendon as in Fig. 6C ***Xanthophysca* Fairmaire**
- Procoxal cavities open posteriorly. Pronotal punctation mostly uniform and sparse (Fig. 1D). Frontal grooves, if distinguishable, short and moderately impressed (Fig. 1E). Hind femur without a distinct emargination on the posterior side. Ductus spermathecae present (Fig. 1G). General shape of the median lobe of the aedeagus clearly flatter (Fig. 1F). Metafemoral extensor tendon as in Fig. 1H..... ***Afrotropicaltica* gen. nov.**
- 5 Eye large and generally distinctly convex (Fig. 5B). Antennae pectinate (generally in males) (Fig. 5A) or clearly serrate. Body distinctly elongate (Fig. 5A), with elytra subparallel laterally, or wider in the apical third. Extended arm of the metafemoral extensor tendon ~ 1/2 the length of the dorsal lobe; apical margin C-shaped (Fig. 5D). Median lobe of the aedeagus (Fig. 5C) straight in lateral view, with apical part thickset and bent ventrally. Spermatheca (Fig. 5E) with basal part roughly subcylindrical, and ductus apically inserted ***Polyclada* Chevrolat**
- Eye small and moderately convex (Fig. 4B). Antennae with subtriangular or moderately serrate middle segments (Fig. 4A). Body generally robust, elliptical, more or less elongate, with elytra rounded laterally (Fig. 4A). Extended arm of the metafemoral extensor tendon distinctly longer than 1/2 the length of the dorsal lobe; apical margin widely arched (am: Fig. 4E). Median lobe of the aedeagus (Fig. 4C) slightly curved in lateral view, with apical part flat, depressed on the ventral surface, and softly curved ventrally. Spermatheca (Fig. 4D) with basal part generally slightly sinuate, dorsally bent at the ductus attachment..... ***Diamphidia* Gerstaecker**

List of *Diamphidia* species

Diamphidia femoralis Gerstaecker, 1855, type species
Diamphidia angolensis Jacoby, 1882
Diamphidia brevis Laboissière, 1942
Diamphidia concinna Weise, 1906
Diamphidia curtula Achard, 1922
Diamphidia flaveola Laboissière, 1941
Diamphidia flaviceps (Allard, 1887)
Diamphidia jacobyi (Gestro, 1895)
Diamphidia lesnei Achard, 1922
Diamphidia limbata (Allard, 1887)
Diamphidia locusta Fairmaire, 1893
Diamphidia longula Weise, 1902
Diamphidia nigrifrons (Allard, 1887)
Diamphidia nigripennis (Jacoby, 1888)
Diamphidia nigroornata Stål, 1858
Diamphidia patricki Bryant, 1948
Diamphidia rhabdoscelis Laboissière, 1942
Diamphidia robusta (Allard, 1887)

Diamphidia rugipennis Laboissière, 1942

Diamphidia semiopaca Achard, 1922

Diamphidia simplex (Peringuey, 1892)

Diamphidia uniformis (Jacoby, 1903)

Diamphidia vittatipennis Baly, 1865




Figure 2. *Blepharidina* Bechyné **A** habitus of *Blepharidina* (*Blepharidina*) *guttulata* (Baly), male, modified from D'Alessandro et al. (2019) **B** *ibid*, head, and pronotum **C** head and pronotum of *Blepharidina* (*Afroblepharida*) *gedyei*, male, Kenya, Sosoma (BAQ) **D** metafemoral extensor tendon, modified from Biondi et al. (2017) **E** head of *Blepharidina intermedia*, modified from Biondi et al. (2017). Abbreviations: am: apical margin; dl: dorsal lobe; ea: extended arm. Scale bars: 3 mm (**A**); 1 mm (**B**, **C**, **E**); 500 µm (**D**).




Figure 3. *Calothea* Heyden **A** habitus of *Calothea haroldi* (Baly), male, Republic of South Africa, Kranskloof (SANC) **B** ibid, head and pronotum **C** head and pronotum of *Calothea marginalis*, modified from Biondi et al. (2017) **D** ibid, metafemoral extensor tendon **E** ibid, head. Abbreviations: am: apical margin; bf: basal furrow; dl: dorsal lobe; ea: extended arm; fg: frontal groove; ps: punctate stria. Scale bars: 3 mm (**A**); 1 mm (**B, C**); 500 µm (**D, E**).

List of *Polyclada* species

- Polyclada pectinicornis* (Olivier, 1791), type species
- Polyclada benti* Gahan, 1895
- Polyclada bohemani* (Baly, 1861)
- Polyclada coriacea* Achard, 1922
- Polyclada kenyensis* Bryant, 1942

Polyclada maculicollis Bryant, 1942
Polyclada ornata (Baly, 1861)
Polyclada ornatipennis Bryant, 1942
Polyclada somaliensis Bryant, 1948
Polyclada variegata (Weise, 1900)




Figure 4. *Diamphidia* Gerstaecker **A** habitus of *Diamphidia femoralis*, male, Mozambique, 25 km N Tete (NMPC) **B** *ibid*, pronotum and base of elytra **C** *ibid*, median lobe of the aedeagus, from left to right in ventral, dorsal, and lateral view, Zambia, Victoria Falls (BAQ) **D** *ibid*, spermatheca, Transvaal, Pretoria (MSNG) **E** *ibid*, metafemoral extensor tendon, Republic of South Africa, Blyde River Canyon (BAQ). Abbreviations: am: apical margin; dl: dorsal lobe; ea: extended arm. Scale bars: 3 mm (**A**); 1 mm (**B**, **C**); 500 µm (**D**, **E**).




Figure 5. *Polyclada* Chevrolat **A** habitus of *Polyclada pectinicornis*, male, modified from Biondi et al. (2022) **B** *ibid*, pronotum and base of elytra, male, Somalia, Bardera (BAQ) **C** *ibid*, median lobe of the aedeagus, from left to right in ventral, dorsal, and lateral view, modified from Biondi et al. (2022) **D** *ibid*, spermatheca **E** *ibid*, metafemoral extensor tendon, Tanzania, Mto. Wa Mbu (BAQ). Abbreviations: am: apical margin; dl: dorsal lobe; ea: extended arm. Scale bars: 3 mm (**A**); 1 mm (**B**, **C**); 500 μ m (**D**, **E**).

List of *Xanthophysca* species

- Xanthophysca perrieri* Fairmaire, 1901, type species
- Xanthophysca androyana* Achard, 1915
- Xanthophysca donckieri* Achard, 1915
- Xanthophysca kolbei* Weise, 1910
- Xanthophysca variegata* Achard, 1915




Figure 6. *Xanthophysca* Fairmaire **A** habitus of *Xanthophysca perrieri*, male, modified from Biondi and D'Alessandro (2012) **B** *ibid*, pronotum and base of elytra **C** metafemoral extensor tendon, Madagascar Ouest, Antsingy (MNHN) **D** *ibid*, median lobe of the aedeagus, from left to right in ventral, dorsal, and lateral view **E** *ibid*, spermatheca, Madagascar, Ankarafantsika Rés. nat. (MNHN). Scale bars: 3 mm (**A**); 1 mm (**B**, **D**); 500 μ m (**C**, **E**).

The taxonomic position at genus level of some species is unclear because they lack one or more characters typical of *Diamphidia*, *Polyclada*, or *Xanthophysca*. These are *Polyclada bedeli* Achard, 1922 with antennae of the *Diamphidia* type; *Polyclada flavipennis* Bryant, 1942 and *P. marginata* Bryant, 1948 with antennae of the *Diamphidia* type and aedeagus lacking the typical *Polyclada* characters; *Polyclada flexuosa* (Baly, 1865) with aedeagus different from the typical *Polyclada* species; *Diamphidia ornaticollis* Bryant, 1948, with closed procoxal cavities and filiform antennae.

Discussion

Afrotropicaltica fulvipennis (Jacoby) comb. nov. was referred to different genera in the previous publications (*Cladocera*, *Blepharida*, *Diamphidia*). However, based on its unique combination of characters, it is not attributable to any of the known flea beetle genera. Considering the diagnostic characters used to identify the main groups of Afrotropical flea beetles (Biondi and D'Alessandro 2012), *Afrotropicaltica* gen. nov. can be grouped with *Diamphidia*, *Blepharidina*, *Calotheca*, *Polyclada*, and *Xanthophysca*, traditionally attributed to the *Blepharida* group sensu Furth and Lee (2000). The following traits characterise these genera: antennae with 11 antennomeres; apical tarsomere of metatarsus simple (not swollen); dorsal margin of middle and hind tibiae with distinct ciliate emargination, which is acute or subrounded apically; prothorax distinctly depressed dorsally (not subcylindrical); first metatarsomere as long as the second or longer, wide, subtriangular or subrounded; body length generally ≥ 4.00 mm; claws generally appendiculate; pronotal sculpture variable but without any ante-basal transverse sulcus. Differences in antennae, pronotum, and elytra are helpful for genera and subgenera identification, as reported in the diagnostic key above.

Genus *Xanthophysca* lacks a comprehensive revision, but the diagnostic characters are coherent among the species attributed to it. Differently, some species currently attributed to *Polyclada* and *Diamphidia* show a combination of characters that makes them not referrable as *Polyclada* or *Diamphidia*. Therefore, *Xanthophysca* and especially *Diamphidia* and *Polyclada* need a revision, also based on the undetermined material and undescribed species present in public and private collections.

Furth and Lee (2000), while providing their morphological synthesis of the *Blepharida* group of genera, stated that some characters are not shared by all the taxa. This is true also for the genera discussed here. Assessing the monophyly of the group and the evolutionary affinity among the included taxa requires a more extensive investigation, also requiring molecular data. The same problems are present for assessing the existence of a wider *Blepharida* group clade.

Acknowledgments

We are grateful to the collection managers and curators who enabled us to study the material preserved in their Institutions: Antoine Mantilleri (MNHN), Roberto Poggi (MSNG), Michael Geiser (NHMUK), Lukas Sekerka (NMPC), and Didier Van den Spiegel and Stéphane Hanot (RMCA). We sincerely thank Alexander S. Konstantinov (National Museum of Natural History, Washington DC, USA) and David G. Furth (Department of Entomology, Smithsonian Institution, Washington DC, USA) for spending time improving our manuscript.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

No funding was reported.

Author contributions

Conceptualization: MB, PD. Data curation: PD. Investigation: PD. Resources: PD, MB. Validation: MB. Visualization: PD, MB. Writing - original draft: PD. Writing - review and editing: MB.

Author ORCIDs

Paola D'Alessandro  <https://orcid.org/0000-0002-4481-9152>

Maurizio Biondi  <https://orcid.org/0000-0003-2190-7376>

Data availability

All of the data that support the findings of this study are available in the main text.

References

- Achard J (1915) Halticides malgaches du genre *Xanthophysca* Fairm. [Col. Phytophaga]. Annales de la Société entomologique de France 84(4): 459–462. <https://doi.org/10.1080/21686351.1915.12279409>
- Achard J (1922) Descriptions de nouveaux Halticini (Col. Chrysomelidae). Fragments Entomologiques 27(3): 3–10. <https://doi.org/10.3406/bsef.1922.26922>
- Allard E (1887) Bulletin entomologique: diagnoses de quelques espèces nouvelles de Coléoptères Phytophages. Annales de la Société entomologique de France 7(6e série): CC–CCI.
- Baly JS (1861) Descriptions of new genera and species of Phytophaga. The Journal of Entomology 1: 193–206, 275–302, 450–459.
- Baly JS (1865) LV.—Descriptions of new genera and species of Gallernucidae. Annals and Magazine of Natural History 16: 402–410. <https://doi.org/10.1080/00222936508679458>
- Becerra JX (2004) Molecular systematics of *Blepharida* beetles (Chrysomelidae: Alticinae) and relatives. Molecular Phylogenetics and Evolution 30: 107–117. [https://doi.org/10.1016/S1055-7903\(03\)00158-1](https://doi.org/10.1016/S1055-7903(03)00158-1)
- Bechyné J (1960) Alticinae (Coleoptera, Phytophagoidea). Exploration du Parc National de l'Upemba. I. Miss. G.F. de Witte 59(3): 39–114.
- Bechyné J (1968) Contribution à la faune du Congo (Brazzaville). Mission A. Villiers et A. Descarpentries. LXXXI. Coleopteres Alticinae. Bulletin de L'institut Francais d'Afrique Noire (série A) 30: 1687–1728.
- Biondi M, D'Alessandro P (2010) Genus-group names of Afrotropical flea beetles (Coleoptera: Chrysomelidae: Alticinae): Annotated catalogue and biogeographical notes. European Journal of Entomology 107: 401–424. <https://doi.org/10.14411/eje.2010.049>
- Biondi M, D'Alessandro P (2012) Afrotropical flea beetle genera: A key to their identification, updated catalogue and biogeographical analysis (Coleoptera, Chrysomelidae, Galerucinae, Alticini). Zookeys 253: 1–158. <https://doi.org/10.3897/zookeys.253.3414>
- Biondi M, Frasca R, Grobbelaar E, D'Alessandro P (2017) Supraspecific taxonomy of the flea beetle genus *Blepharida* Chevrolat, 1836 (Coleoptera: Chrysomelidae) in the Afrotropical Region and description of *Afroblepharida* subgen. nov. Insect Systematics & Evolution 48: 97–155 <https://doi.org/10.1163/1876312X-48022152>

- Biondi M, Iannella M, D'Alessandro P (2019) Unravelling the taxonomic assessment of an interesting new species from Socotra Island: *Blepharidina socotrana* sp. nov. (Coleoptera: Chrysomelidae). *Acta Entomologica Musei Nationalis Pragae* 59(2): 499–505. <https://doi.org/10.2478/aemnp-2019-0040>
- Biondi M, D'Alessandro P, Cerasoli F, De Simone W, Iannella M (2022) Taxonomy, Habitat Preference, and Niche Overlap of Two Arrow-Poison Flea Beetle Species of the Genus *Polyclada* in Sub-Saharan Africa (Coleoptera, Chrysomelidae). *Insects* 13: 668. <https://doi.org/10.3390/insects13080668>
- Biondi M, Iannella M, D'Alessandro P (2024) Ecological profile of the flea beetle genus *Calotheca* Heyden in South Africa (Chrysomelidae, Galerucinae, Alticinae). *Insects* 15: 994. <https://doi.org/10.3390/insects15120994>
- Bryant GE (1942) New species of *Polyclada* (Coleoptera, Halticinae) from Africa. *Proceedings of the Royal Entomological Society of London. Series B, Taxonomy* 11(11): 161–165. <https://doi.org/10.1111/j.1365-3113.1942.tb00716.x>
- Bryant GE (1948) XII.—New species of Chrysomelidae (Halticinae) from East Africa, *Annals and Magazine of Natural History, Series 12*, 1(3): 145–171. <https://doi.org/10.1080/00222934808653898>
- Calcetas OA, Staines CI (2024) New Species of the Genus *Podontia* Dalman (Coleoptera: Chrysomelidae: Galerucinae: Alticini) in the Philippines and Redescription of *Podontia flava* Baly from Sarawak. *Philippine Journal of Science* 153 (3): 903–909. <https://doi.org/10.56899/153.03.13>
- Chaboo CS, Grobbelaar E, Larsen A (2007) Fecal Ecology in Leaf Beetles: Novel Records in the African Arrow-Poison Beetles, *Diamphidia* Gerstaecker and *Polyclada* Chevrolat (Chrysomelidae: Galerucinae). *The Coleopterist Bulletin* 61: 297–309. [https://doi.org/10.1649/0010-065X\(2007\)61\[297:FEILBN\]2.0.CO;2](https://doi.org/10.1649/0010-065X(2007)61[297:FEILBN]2.0.CO;2)
- Chapuis MF (1875) *Histoire naturelle des Insectes. Genera des Coléoptères ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'Insectes*. Par M. M. Lacordaire et F. Chapuis. Tome onzième. Famille des Phytophages. Paris, 420 pp. [+ pls 124–134]
- Chevrolat (1837) *Catalogue des Coléoptères de la collection de M. le Comte Dejean*. Troisième édition, revue, corrigée et augmentée, livr. 5. Dejean PFMA (Ed.). Librairie Méquignon-Marvis Père et Fils, Paris, [i–xiv +] 385–503]. <https://www.biodiversitylibrary.org/item/285852#page/390/mode/1up>
- D'Alessandro P, Biondi M (2023) Generic placement of the African flea beetle *Polyclada maculipennis* Bryant and the possible occurrence of the genus *Procalus* Clark in sub-Saharan Africa (Coleoptera, Chrysomelidae, Galerucinae, Alticini). *ZooKeys* 1145: 181–189. <https://doi.org/10.3897/zookeys.1145.90667>
- D'Alessandro P, Samuelson A, Biondi M (2016) Taxonomic revision of the genus *Arsipoda* Erichson, 1842 (Coleoptera, Chrysomelidae) in New Caledonia. *European Journal of Taxonomy* 230: 1–61. <https://doi.org/10.5852/ejt.2016.230>
- D'Alessandro P, Frasca R, Grobbelaar E, Iannella M, Biondi M (2018a) Systematics and biogeography of the Afrotropical flea beetle subgenus *Blepharidina* (*Afroblepharida*) Biondi & D'Alessandro, with description of seven new species (Coleoptera, Chrysomelidae, Galerucinae, Alticini). *Insect Systematics & Evolution* 49: 443–480. <https://doi.org/10.1163/1876312X-00002182>
- D'Alessandro P, Iannella M, Frasca R, Biondi M (2018b) Distribution patterns and habitat preference for the genera-group *Blepharida* s.l. in Sub-Saharan Africa (Coleoptera: Chrysomelidae: Galerucinae: Alticini). *Zoologischer Anzeiger* 277: 23–32. <https://doi.org/10.1016/j.jcz.2018.08.001>

- D'Alessandro P, Iannella M, Biondi M (2019) Revision of the Afrotropical flea beetle subgenus *Blepharidina* s. str. Bechyné (Coleoptera, Chrysomelidae). *Zootaxa* 4545(1): 032–060. <https://doi.org/10.11646/zootaxa.4545.1.2>
- D'Alessandro P, Iannella M, Grobbelaar E, Biondi M (2020) Revision of the *Calotheca nigrotessellata* species group from southern Africa, with description of two new species (Coleoptera: Chrysomelidae: Galerucinae: Alticini). *Fragmenta entomologica* 52(2): 169–182. <https://doi.org/10.4081/fe.2020.457>
- D'Alessandro P, Iannella M, Grobbelaar E, Biondi M (2021) Taxonomic revision of the *Calotheca parvula* species group from southern Africa, with descriptions of three new species (Coleoptera, Chrysomelidae). *African Invertebrates* 62(1): 315–337. <https://doi.org/10.3897/AfrInvertebr.62.62426>
- D'Alessandro P, Iannella M, Grobbelaar E, Biondi M (2022) *Calotheca nigromaculata* species-group from sub-Saharan Africa with descriptions of two new species from Kwa-Zulu-Natal (Chrysomelidae, Galerucinae, Alticini). *ZooKeys* 1084: 119–137. <https://doi.org/10.3897/zookeys.1084.73175>
- D'Alessandro P, Grobbelaar E, Iannella M, Biondi M (2023a) *Calotheca borowieci* sp. nov. and *C. orophila* sp. nov. from southern Africa (Coleoptera, Chrysomelidae, Galerucinae, Alticini). *Annales Zoologici* 73 (3): 421–428. <https://doi.org/10.3161/00034541ANZ2023.73.3.006>
- D'Alessandro P, Grobbelaar E, Iannella M, Biondi M (2023b) *Calotheca leonardii*, a new flea beetle species from southern Africa (Coleoptera: Chrysomelidae: Galerucinae: Alticini) *Natural History Sciences. Atti della Società italiana di Scienze naturali del Museo civico di Storia naturale di Milano* 10(Supplement 1): 5–10. <https://doi.org/10.4081/nhs.2023.681>
- Döberl M (1986) Die Spermathek als Bestimmungshilfe bei den Alticinen. *Entomologische Blätter für Biologie und Systematik der Käfer* 82(1–2): 3–14.
- Evenhuis NL (2021) The insect and spider collections of the world website. <http://hbs.bishopmuseum.org/codens/> [Accessed 09 October 2024]
- Fairmaire L (1893) Description d'une nouvelle espèce de Chrysomélide de l'Afrique australe. *Annales de la Société entomologique de France* 62. Bulletin entomologique: CCCXLVII.
- Fairmaire L (1901) Matériaux pour la faune coléoptérique de la région malgache (11^e note) (fin). *Revue d'Entomologie* 20: 101–248.
- Furth DG (1982) The Metafemoral Spring of Flea Beetles (Chrysomelidae: Alticinae). *Spixiana* 7: 11–27.
- Furth DG (1992) The New World *Blepharida* group, with a key to genera and description of a new species (Coleoptera: Chrysomelidae). *Journal of New York Entomological Society* 100: 399–414. <http://www.jstor.org/stable/25009970>
- Furth DG (1998) New World *Blepharida* Chevrolat 1836 (Coleoptera: Chrysomelidae: Alticinae). *Memoirs of the Entomological Society of Washington* 21: 1–109.
- Furth DG, Lee JE (2000) Similarity of the *Blepharida*-group genera using larval and adult characters (Coleoptera: Chrysomelidae: Alticinae). *Journal of the New York Entomological Society* 108: 26–51. [https://doi.org/10.1664/0028-7199\(2000\)108\[0026:SOT-BGG\]2.0.CO;2](https://doi.org/10.1664/0028-7199(2000)108[0026:SOT-BGG]2.0.CO;2)
- Furth DG, Young DA (1988) Relationships of herbivore feeding and plant flavonoids (Coleoptera: Chrysomelidae and Anacardiaceae: *Rhus*). *Oecologia* 74: 496–500. <https://doi.org/10.1007/BF00380045>
- Gahan (1895) On the Coleoptera obtained by Dr. Anderson's Collector during Mr. T. Bent's Expedition to the Hadramaut, South Arabia. *The Journal of the Linnean Society of London. Zoology* 25(161): 285–291. <https://doi.org/10.1111/j.1096-3642.1895.tb03418.x>

- Gerstaecker CEA (1855) Diagnosen der von Peters in Mossambique Gesammelten neuer Coleopteren und Lepidopteren. Monatsberichte der Akademie der Wissenschaften zu Berlin 1855: 636–639.
- Gestro R (1895) Esplorazione del Giuba e dei suoi affluenti compiuta dal Cap. V. Bottego durante gli anni 1892-93 sotto gli auspici della Società Geografica Italiana. Risultati Zoologici. XVI. Coleotteri. Anali del Museo Civico di Storia Naturale di Genova, serie 2a, 15(35): 249–478. <https://doi.org/10.5962/bhl.title.49163>
- Heyden von L (1887) Kleine coleopterologische Mitteilungen. Wiener Entomologische Zeitung 6: e98. <https://doi.org/10.5962/bhl.part.17738>
- Hope FW (1840) The coleopterist's manual, part the third, containing various families, genera, and species, of beetles, recorded by Linneus and Fabricius. Also, descriptions of newly discovered and unpublished insects. Bowdery & Kerby, London, 191 pp.
- Iannella M, D'Alessandro P, De Simone W, Biondi M (2021) Habitat Specificity, Host Plants and Areas of Endemism for the Genera-Group *Blepharida* s.l. in the Afrotropical Region (Coleoptera, Chrysomelidae, Galerucinae, Alticini). Insects 12: 299. <https://doi.org/10.3390/insects12040299>
- Jacoby M (1882) Descriptions of new genera and species of phytophagous Coleoptera. Proceedings of the Zoological Society of London 50(1): 50–58. <https://doi.org/10.1111/j.1096-3642.1882.tb02727.x>
- Jacoby M (1888) Descriptions of new or little-known species of phytophagous Coleoptera from Africa and Madagascar. Transactions of the Entomological Society of London 189–206. <https://doi.org/10.1111/j.1365-2311.1888.tb01308.x>
- Jacoby M (1895) 179. Chrysomeliden von Togo (Bismarckburg). Deutsche Entomologische Zeitschrift, 165–188.
- Jacoby M (1903) Descriptions of new genera and species of Phytophagous Coleoptera. Stettiner Entomologische Zeitung 64: 292–336.
- Koch C (1958) Preliminary notes on the coleopterological aspect of the arrow poison of the bushmen. Pamphlet of the South African Biological Society 20: 49–54.
- Kolbe HJ (1894) Der Pfeilgiftkäfer der Kalahari-Wüste, *Diamphidia simplex* Péring. (*Locusta* Fairm.). Entomologische Zeitung 55: 79–86.
- Laboissière V (1941) Observations sur les Halticinae de la Collection du Musée du Congo et descriptions d'espèces nouvelles. Revue de Zoologie et de Botanique Africaines 34(2): 236–263.
- Laboissière V (1942) Halticinae (Coleoptera, Phytophaga) Fam. Chrysomelidae. Exploration du Parc National Albert. Mission G.F. De Witte (1933–1935). Institut des Parcs Nationaux du Congo Belge 39: 1–131.
- Lee CF, Cheng HT (2007) The Chrysomelidae of Taiwan Vol. 1. Sishou-Hills Insect Observation Network Press, Taipei, 199 pp.
- Medvedev LN (1999) A revision of the group Blepharidiini (Chrysomelidae: Alticinae) from the Oriental region. Russian Entomological Journal 8: 175–184.
- Nadein KS, Betz O (2016) Jumping mechanisms and performance in beetles. I. Flea beetles (Coleoptera: Chrysomelidae: Alticini). Journal of Experimental Biology 219: 2015–2027. <https://doi.org/10.1242/jeb.140533>
- Olivier M (1791) Histoire Naturelle. Insectes. Tome VI. In: Encyclopédie Méthodique, Panckoucke, Paris, France, 368 pp.
- Peringuey L (1892) On beetles collected in Tropical South-Western Africa. In: Peringuey L (Ed.) Third contribution to the South African Coleopterous fauna. Transactions of the South African Philosophical Society 6(2): 1–94. <https://doi.org/10.1080/21560382.1889.9526255>

- Prathapan KD, Chaboo CS (2011) Biology of *Blepharida*-group flea beetles with first notes on natural history of *Podontia congregata* Baly, 1865 an endemic flea beetle from southern India (Coleoptera, Chrysomelidae, Galerucinae, Alticini). Zookeys 157: 95–130. <https://doi.org/10.3897/zookeys.157.1472>
- Scherer G (1962) Beitrag zur kenntnis der Alticiden-fauna Zentral-Afrikas (Coleoptera Chrysomelidae Alticinae). Annales du Musée Royale de l'Afrique Centrale, Tervuren (Ser. 8, Zool.) 113: 1–82.
- Scherer G (1969) Contributions à la connaissance de la faune entomologique de la Côte d'Ivoire (L. Decelle, 1961–1964) XLIII. Coleoptera Chrysomelidae Alticinae. Annales du Musée Royale de l'Afrique Centrale, Tervuren (Ser. 8, Zool.) 175: 365–371.
- Scherer G (1972) Coleoptera aus Nordostafrika. Chrysomelidae: Alticinae. Notulae Entomologicae 52: 1–17.
- Schmitt M, Uhl G (2015) Functional morphology of the copulatory organs of a reed beetle and a shining leaf beetle (Coleoptera: Chrysomelidae: Donaciinae, Criocerinae) using X-ray micro-computed tomography. In: Jolivet P, Santiago-Blay J, Schmitt M (Eds) Research on Chrysomelidae 5. ZooKeys 547: 193–203. <https://doi.org/10.3897/zookeys.547.7143>
- Stål C (1858) Entomologiska bidrag. Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar, 247–258.
- Takizawa H (2005) Supra-generic subdivisions of the subfamily Alticinae based on larval characters, with descriptions of larvae of Hispaniolan species (Coleoptera: Chrysomelidae). Insecta Matsumurana 62: 187–206.
- Weise J (1900) Beschreibungen africanischer Chrysomeliden nebst synonymischen Bemerkungen. Deutsche Entomologische Zeitschrift 2: 446–459. <https://doi.org/10.1002/mmnd.48019000239>
- Weise J (1902) Afrikanische Chrysomeliden. Archiv für Naturgeschichte 68: 119–174.
- Weise J (1906) Ostafrikanische Chrysomeliden und Coccinelliden. Deutsche entomologische Zeitschrift 1: 35–64.
- Weise J (1910) Chrysomelidae von Madagaskar, der Comoren und der Inseln Ostafrikas. In: Reise in Ostafrika in den Jahren 1903–1905 mit Mitteln der Hermann und Elise geb. Heckmann Wentzel-Stiftung ausgeführt. Von Professor Dr. Alfred Voeltzkow. Wissenschaftliche Ergebnisse. Band II. Systematische Arbeiten. Heft V, 419–506.